

Comparison of Ego Strength, Aggression, and Fear of Loss in Three Groups with Anxiety Symptoms, Depression Symptoms, and Normal Individuals

Zahra. Razaghi¹, Atefeh. Hojjati^{2*}

¹ M.A., Student, Department of Psychology, Qom Medical Sciences Branch, Islamic Azad University, Qom, Iran

² Assistant Professor, Department of Psychology, Qom Medical Sciences Branch, Islamic Azad University, Qom, Iran

* Corresponding author email address: hojjati14@yahoo.com

Article Info

Article type:

Original Research

How to cite this article:

Razaghi, Z., & Hojjati, A. (2025). Comparison of Ego Strength, Aggression, and Fear of Loss in Three Groups with Anxiety Symptoms, Depression Symptoms, and Normal Individuals. *Journal of Assessment and Research in Applied Counseling*, 7(4), 1-9.

<http://dx.doi.org/10.61838/kman.jarac.3081>



© 2025 the authors. Published by KMAN Publication Inc. (KMANPUB), Ontario, Canada. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) License.

ABSTRACT

Objective: This study was conducted with the aim of comparing ego strength, aggression, and fear of loss among three groups: individuals with anxiety symptoms, individuals with depression symptoms, and normal individuals.

Methods and Materials: The research design was a causal-comparative method. The statistical population of this study included all women aged 20-40 years who referred to counseling centers in the province of Qom. The sample consisted of 105 women aged 20-40 years who referred to counseling centers in the province of Qom and were selected using convenience sampling. The instruments used in this study included the Beck Depression Inventory (second edition), the Beck Anxiety Inventory, the Ego Strength Psychological Questionnaire by Markstrom et al., the Fear of Loss Questionnaire by Przybylski, and the Buss-Perry Aggression Questionnaire. To analyze the findings, both univariate and multivariate analysis of variance (ANOVA) were used, and data analysis was conducted using SPSS.26 software.

Findings: The results showed that the normal group had higher ego strength compared to the groups with depression and anxiety symptoms. The group with depression symptoms exhibited a higher fear of loss compared to the anxiety symptoms group and the normal group, and aggression was higher in the anxiety symptoms group compared to the depression symptoms group and the normal group ($p < .01$).

Conclusion: The study concludes that ego strength, aggression, and fear of loss significantly differ among individuals with anxiety symptoms, depression symptoms, and normal individuals. Higher ego strength was observed in the normal group, while the depression group exhibited greater fear of loss, and aggression was most pronounced in the anxiety group.

Keywords: Ego Strength, Aggression, Fear of Loss, Anxiety, Depression.

1. Introduction

Anxiety disorder, as defined in the DSM-5, is a mental state characterized by intense arousal, with its primary features being excessive fear, doubt, and worry (Rapee et al., 2023). Cognitive processing in anxiety disorder mirrors that observed in trait anxiety; therefore, understanding the cognitive processes and mechanisms underlying this disorder enhances our general understanding of vulnerability to anxiety (Tam et al., 2023).

In addition to anxiety, all individuals experience depression at some point in their lives. Depression is the most common mental disorder, and its prevalence has been increasing recently (Daly et al., 2021). Depression is generally recognized as sadness and can occur following any unpleasant event (Harsanyi et al., 2022). Depression is a disorder primarily characterized by mood alterations, which are accompanied by distinct changes in behavior, attitude, thinking, efficiency, and physiological functions. As a symptom, depression may be present in various psychological and physical disorders, constituting a secondary component of the clinical presentation (Hull et al., 2021).

Furthermore, depression can be a natural emotion, characterized by mourning and grief. The illness of depression affects the entire organism and impacts all areas of an individual's life (Renaud-Charest et al., 2021). This effect is evident in feelings, energy, motivation, thinking, physical actions, personality, and interests. Depression is an emotional challenge that can be distressing for both the individual and those close to them, often manifesting as persistent boredom (Moncrieff et al., 2023). This condition persists for most of the day and may last for several weeks or longer. The presence of depression can lead to stress, affecting work or social performance. Boredom is often accompanied by sadness or a sense of emptiness (Cunningham et al., 2021).

Various approaches in psychology have examined how individuals respond to unpleasant psychological conditions, with one of the most profound being the psychoanalytic approach (Toprak et al., 2022). According to this approach, personality consists of three components: the id, ego, and superego. Since the ego is responsible for managing the psychic system, psychological problems arise when the ego fails to fulfill its duties (Sadafi et al., 2020). Ego strength refers to the ability to maintain one's identity despite psychological stress, suffering, and the conflict between internal needs and external demands. In other words, it is the

ability to preserve ego stability based on a relatively stable set of personality traits, which reflects in maintaining mental health (Aldahadha Al-Khawaldeh & Mohammed Al-Khawaldeh, 2023). Ego strength essentially indicates an individual's capacity to endure stress without experiencing debilitating anxiety (Toprak et al., 2022). Therefore, it can be said that the strength and resilience of the ego, as well as how defense mechanisms are employed in the face of events and during the experience of unpleasant emotions, are among the factors related to self-injurious behavior (Hafer, 2020).

One variable that appears to influence depression and anxiety disorders is aggression. Aggression is behavior intended to harm another (Kjærviik & Bushman, 2021). Aggressive behaviors manifest in various forms, with the most prominent being physical and verbal aggression. Physical aggression includes activities such as hitting, pushing, shoving, throwing objects, and threatening to perform these actions, all intended to harm others (Falla et al., 2023). Physical aggression, like verbal aggression, is observable through its manifestations and is therefore considered a direct expression of aggression (Wei et al., 2023).

Another component that seems to be present in individuals with depression and anxiety disorders is the fear of missing out (FoMO). FoMO, added as a new term to the dictionary in 2013, is characterized by a pervasive fear that others may be having valuable experiences from which one is absent. This fear is marked by a desire to stay connected with what others are doing, constantly checking social networks, and fearing missing out on events (Cheng & Wee, 2023). Additionally, studies have shown that FoMO is a psychological construct defined by the fear of not being present in rewarding experiences that others are having, coupled with the desire for continuous connection with others' experiences (Radomsky, 2022). This phenomenon, prevalent among most people, can cause significant stress and lead to harmful outcomes (Wisse et al., 2019). Recent studies have linked internet and social media addiction to psychological factors, including increased stress levels, anxiety, depression, lower self-esteem, reduced relationship quality, poor sleep quality, and heightened suicidal thoughts and behaviors, along with reduced life satisfaction among young people (Wei et al., 2022). On the other hand, problematic social media use and its connection with rumination, repetitive thoughts, negative feelings, and loneliness have been observed (Kelly et al., 2016). Based on the aforementioned points, the present study aims to compare ego strength, aggression, and fear of loss in

individuals with anxiety symptoms, depression symptoms, and normal individuals.

2. Methods and Materials

2.1. Study Design and Participants

The present study employed a causal-comparative design. The statistical population consisted of all women aged 20-40 years who referred to counseling centers in Qom province, and their anxiety and depression symptoms were assessed using the Beck Anxiety Inventory (BAI) and the Beck Depression Inventory-II (BDI-II). They were categorized into three groups: those with anxiety symptoms, those with depression symptoms, and normal individuals. The research sample comprised 105 women aged 20-40 years who visited counseling centers in Qom province. They were assessed using the BAI and BDI-II and categorized into three groups: those with anxiety symptoms, those with depression symptoms, and normal individuals. According to Delavar (2012), the minimum sample size in each group for causal-comparative studies is 30 individuals. In this study, using Cohen's formula, considering the number of variables under investigation, an effect size of 1, a test power of 0.8, and a confidence level of 95%, it was estimated that 35 participants per group (35 in the anxiety disorder group, 35 in the depression group, and 35 in the normal group) were required, resulting in a total of 105 participants. The sampling method used in this study was convenience sampling. Inclusion criteria included informed consent to participate in the study, complete filling out of the questionnaires, age between 20 and 40 years, at least elementary education, no substance abuse, willingness to participate in the study, no use of antidepressant or anti-anxiety medications, and no depression or anxiety symptoms for normal individuals. Exclusion criteria included psychosis, the presence of other psychological disorders based on DSM-5-TR, alcohol and substance abuse, incomplete questionnaires, and the use of antidepressant and anti-anxiety medications.

After obtaining the necessary approvals and identifying the participants through convenience sampling at counseling centers in Qom, individuals were screened for anxiety and depression symptoms using the Beck Anxiety Inventory and Beck Depression Inventory-II. For the depression group, individuals with the highest scores on the Beck Depression Inventory-II were selected, and for the anxiety group, those with the highest scores on the Beck Anxiety Inventory were chosen. For the normal group, individuals with normal

scores on both inventories were selected. The three groups—individuals with anxiety symptoms, depression symptoms, and normal individuals in Qom—completed the questionnaires. Participants were instructed to ask the researcher for clarification if they encountered any ambiguities while completing the questionnaires. At the end, participants were thanked for their cooperation. Before the study began, ethical approval was obtained from the university's ethics committee and relevant authorities at the research site. Participants were provided with written information about the study, assured of the confidentiality of their information, informed that the data would be used solely for research purposes, and assured that their participation was voluntary. Their names and personal information were not recorded to protect their privacy, and written informed consent was obtained from all participants.

2.2. Measures

2.2.1. Depression

The Beck Depression Inventory was designed and developed by Beck to measure the severity of depression in adults diagnosed with depressive disorders. The inventory consists of 21 items, rated on a four-point Likert scale, with items such as "I feel that there is no hope for the future and that things will only get worse." The inventory is self-rated. In this study, depression is defined by the score obtained by respondents on the 21-item Beck Depression Inventory. The inventory is unidimensional, and a cut-off score of 18 can correctly identify up to 92% of patients with major depressive disorder. The following ranges indicate the severity of depression: 0 to 13: minimal or no depression; 14 to 19: mild depression; 20 to 28: moderate depression; and 29 to 63: severe depression. The Cronbach's alpha coefficient reported in the study by Taheri Tanjani (2014) for this inventory was above 0.7 (Bahmani Nia & Sohrabi Shegefti, 2024). The reliability of this inventory, using Cronbach's alpha, was reported as 0.79.

2.2.2. Fear of Missing Out

To measure the Fear of Missing Out (FoMO), the questionnaire developed by Przybylski (2013) was used. It consists of 10 items rated on a five-point Likert scale (not at all true, slightly true, moderately true, mostly true, extremely true). Scores range from 1 (not at all true) to 5 (extremely true), with total scores ranging from 10 to 50. A score above 20 indicates a higher level of FoMO. This questionnaire was

translated into Persian and validated by Bayrami and colleagues in 2019, with a Cronbach's alpha of 0.87 (Karami Boldaghi & Daryazadeh, 2018). The reliability of this questionnaire, using Cronbach's alpha, was reported as 0.79.

2.2.3. Aggression

The Buss-Perry Aggression Questionnaire, a revised version of the earlier Hostility Questionnaire, was developed by Buss and Perry (1992). It is a self-report tool comprising 29 items and four subscales: physical aggression (PA), verbal aggression (VA), anger (A), and hostility (H). Respondents rate each item on a five-point scale from "very much like me" (5) to "not at all like me" (1). Items 9 and 16 are reverse-scored. The total aggression score is obtained by summing the subscale scores. The questionnaire has demonstrated acceptable validity and reliability, with test-retest coefficients for the four subscales (over a 9-week interval) ranging from 0.72 to 0.80, and intercorrelations among the subscales ranging from 0.38 to 0.49. Internal consistency, as measured by Cronbach's alpha, was 0.82 for physical aggression, 0.81 for verbal aggression, 0.83 for anger, and 0.80 for hostility (Adavi et al., 2017; Ramazani & Hadizadeh Kafash, 2016). The reliability of this questionnaire, using Cronbach's alpha, was reported as 0.79, 0.77, 0.78, and 0.76, respectively.

2.2.4. Anxiety

Assessing anxiety symptoms is crucial in diagnosis and treatment. Although various scales have been developed based on different perspectives, these scales have been criticized for theoretical conceptualization and methodological issues. To address these issues, Aaron Beck and his colleagues (1990) developed the Beck Anxiety Inventory (BAI), specifically designed to measure the severity of clinical anxiety symptoms in individuals. The BAI is a self-report questionnaire for measuring anxiety severity in adolescents and adults. The total score ranges from 0 to 63. The internal consistency (alpha coefficient) of the BAI is 0.92, its one-week test-retest reliability is 0.75, and the item-total correlations range from 0.30 to 0.76. Five types of validity—content, concurrent, construct, diagnostic, and complete—have been assessed, all indicating the BAI's

high efficacy in measuring anxiety severity. Kaviani and Mousavi (2008) reported a validity coefficient of approximately 0.72, a one-month test-retest reliability of 0.83, and a Cronbach's alpha of 0.92 for the BAI in the Iranian population (Soheili-Pour et al., 2023).

2.2.5. Psychological Ego Strength

This inventory was developed by Markstrom, Sabino, Turner, and Berman in 1997 to assess ego strength. It consists of 64 items and 8 subscales: ego strength, hope, will, purpose, competence, fidelity, love, care, and wisdom. The inventory uses a five-point Likert scale (not at all like me, somewhat unlike me, no opinion, somewhat like me, very much like me), with scores of 1-2-3-4-5 assigned accordingly. The score for each subscale is obtained by summing the scores of the relevant items. The highest possible score is 320, and the lowest is 64. A score close to 320 indicates a high level of ego strength, while a score close to 64 indicates low ego strength. Markstrom and colleagues (1997) confirmed the face, content, and construct validity of the inventory. They reported a Cronbach's alpha of 0.68 for reliability. Altafi (2009) reported a Cronbach's alpha of 0.91 and a split-half reliability of 0.77 for the inventory in an Iranian sample (Zanjanchi Niko & Farahani, 2024). The reliability of this inventory, using Cronbach's alpha, was reported as 0.76.

2.3. Data analysis

For data analysis in this study, the following statistical methods were used: descriptive statistics, including mean, standard deviation, skewness, and kurtosis. For inferential statistics, univariate and multivariate analysis of variance (ANOVA) was used, and data analysis was conducted using SPSS version 26.

3. Findings and Results

The demographic characteristics of the study sample are presented with frequency and percentage. As shown, each of the three studied groups consisted of 35 participants, representing 33.33% of the total sample for each group.

Table 1

Descriptive Statistics of the Studied Groups

Variable	Group	Mean	Standard Deviation	N
Ego Strength	Anxiety	156.94	64.15	35
	Depression	114.97	42.09	35
	Normal	201.49	56.93	35
Fear of Loss	Anxiety	30.46	10.67	35
	Depression	38.49	9.11	35
	Normal	21.09	7.58	35
Aggression	Anxiety	80.51	26.13	35
	Depression	63.09	31.88	35
	Normal	39.37	24.84	35

Table 1 shows the descriptive statistics for the studied groups based on the variables under investigation, including mean and standard deviation for each group. The normal group had a higher ego strength score than the other groups. Additionally, the anxiety group scored higher on ego strength compared to the depression group. The depression group had a higher fear of loss score compared to the anxiety group. Aggression was higher in the anxiety group. Data from all 105 observations were analyzed, with no missing or lost data found. Univariate outliers were identified using frequency tables and box plots, with no outliers observed. Therefore, all 105 observations were ready for analysis. The box plot below illustrates the overall variables of the study.

The results also indicated that for all variables and their subscales, the skewness and kurtosis divided by the standard error fell within the range of -2 to +2, suggesting that the distribution of scores in the three groups was not non-normal, meaning this assumption was met. Box's M test examines the null hypothesis that the observed covariance matrices of the dependent variables are equal across groups. Since the value of F (1.224) is not significant at the given error level ($p = .259$), the null hypothesis is not rejected. This means that the observed covariance matrices between the different groups are equal.

Table 2

Results of Multivariate Tests

Effect	Value	F	Hypothesis df	Error df	Sig	Partial Eta Squared
Pillai's Trace	.766	20.916	6	202	.001	.383
Wilks' Lambda	.350	23.010	6	200	.001	.408
Hotelling's Trace	1.524	25.154	6	198	.001	.433
Roy's Largest Root	1.261	42.443	3	101	.001	.558

Table 2 reports the results of Pillai's Trace, Wilks' Lambda, Hotelling's Trace, and Roy's Largest Root tests. All four tests are used to assess the significance of multivariate analysis of variance (MANOVA). As noted, Wilks' Lambda is the most commonly reported test, while Pillai's Trace is considered the most robust when assumptions are violated. When all conditions are met and sample sizes are equal, the

results of all four tests tend to be similar. The results of the Pillai's Trace test (Sig = .000), Wilks' Lambda test (Sig = .000), Hotelling's Trace test (Sig = .000), and Roy's Largest Root test (Sig = .000), all with p-values less than .05, indicate that there is a significant difference between the three groups in at least one of the variables or components examined.

Table 3

Results of Variance Analysis

Source	Dependent Variable	Sum of Squares	df	Mean Square	F	Sig	Partial Eta Squared
Group	Ego Strength	191997.390	2	95998.695	31.554	.000	.382
	Fear of Loss	5308.819	2	2654.410	31.314	.000	.380
	Aggression	29853.333	2	14926.667	19.330	.000	.275
Error	Ego Strength	310320.171	102	3042.355			
	Fear of Loss	8646.171	102	84.766			
	Aggression	78763.657	102	772.193			
Total	Ego Strength	2923376.000	105				
	Fear of Loss	108515.000	105				
	Aggression	499200.000	105				

Table 3 presents the final results of the analysis of variance (ANOVA). To test the hypotheses and examine the differences in means between the groups, the significance

level of the test should be referred to. The results indicate significant differences between the three groups in ego strength, fear of loss, and aggression.

Table 4

Results of Bonferroni Post Hoc Test for Ego Strength

Variable	(I) Group	(J) Group	Mean Difference (I-J)	Standard Error	Sig	Lower Bound (95% CI)	Upper Bound (95% CI)
Ego Strength	Anxiety	Depression	59.82857*	13.18517	.000	27.7345	91.9227
		Normal	-44.54286*	13.18517	.003	-76.6369	-12.4488
	Depression	Anxiety	-59.82857*	13.18517	.000	-91.9227	-27.7345
		Normal	-104.37143*	13.18517	.000	-136.4655	-72.2773
	Normal	Anxiety	44.54286*	13.18517	.003	12.4488	76.6369
		Depression	104.37143*	13.18517	.000	72.2773	136.4655
Fear of Loss	Anxiety	Depression	-8.02857	2.20086	.001	-13.3857	-2.6714
		Normal	9.37143	2.20086	.000	4.0143	14.7286
	Depression	Anxiety	8.02857	2.20086	.001	2.6714	13.3857
		Normal	17.40000	2.20086	.000	12.0429	22.7571
	Normal	Anxiety	-9.37143	2.20086	.000	-14.7286	-4.0143
		Depression	-17.40000	2.20086	.000	-22.7571	-12.0429
Aggression	Anxiety	Depression	17.42857	6.64269	.030	1.2596	33.5976
		Normal	41.14286	6.64269	.000	24.9739	57.3118
	Depression	Anxiety	-17.42857	6.64269	.030	-33.5976	-1.2596
		Normal	23.71429	6.64269	.002	7.5453	39.8833
	Normal	Anxiety	-41.14286	6.64269	.000	-57.3118	-24.9739
		Depression	-23.71429	6.64269	.002	-39.8833	-7.5453

As the p-value for the Bonferroni post hoc test was estimated to be less than .05, it can be concluded that there is a significant difference between the three groups for each of the variables: ego strength, fear of loss, and aggression (Table 4).

4. Discussion and Conclusion

Based on the results obtained, it is evident that the normal group had higher ego strength compared to the depression and anxiety groups. The depression group exhibited a higher

mean fear of loss compared to the anxiety and normal groups. There is a significant difference in aggression among the normal, anxiety, and depression groups. Comparing the three groups showed that the aggression variable score was lower in the normal group compared to the anxiety and depression groups, and higher in the anxiety group compared to the depression group. These findings are consistent with prior studies (Bahmani Nia & Sohrabi Shegefti, 2024; Madresi & Shomali Askuei, 2023; Malhotra et al., 2023; Nikosafat & Qarebaghi, 2020; Tao et al., 2023; Zanjanchi Niko & Farahani, 2024).

In explaining these findings, it can be said that the ego strives to balance external reality and internal pressures. Ego strength is a strong predictor of success in facing life's difficult situations, and weakness in the ego leads to resorting to ineffective strategies. From a mental health perspective, ego strength is the ability of individuals to maintain identity and a sense of self when confronted with pain, distress, and conflicts. A strong ego has the capacity to develop new defense mechanisms and coping strategies. Ego strength is related to our core sense of self and is a part of our psychological, social, emotional, and cultural growth, indicating the ability to adapt and be flexible in responding to life's challenging situations. This ability influences an individual's capacity to regulate difficult emotions, accept and endure stress, discomfort, and disappointment without losing emotional control. On the other hand, a deficiency in ego strength can lead to poor judgment, difficulty in reality assessment, and interpersonal problems (Tao et al., 2023). Low ego strength is associated with depression symptoms such as guilt and the lack of impact on the environment. Therefore, a strong ego allows individuals to exhibit fewer psychological distress symptoms and to show more tolerance and resilience in the face of life's hardships.

Another variable with a significant difference among the three groups is aggression. Comparing the three groups revealed that the aggression variable score was lower in the normal group compared to the anxiety and depression groups, and higher in the anxiety group compared to the depression group. Aggression is an antisocial behavior characterized by harming others directly or indirectly (Adavi et al., 2017). Uncontrolled aggressive behavior not only leads to interpersonal problems and violations of others' rights but may also be internalized, causing various physical-psychological problems. Research indicates that mental rumination or revisiting events significantly increases anger and aggression, which may manifest not only in physical but also in verbal behaviors (Kinrade et al., 2022).

In explaining this finding, it can be said that aggression differs among individuals with anxiety, depression, and normal symptoms. Individuals with anxiety symptoms may seek experiences that demonstrate stress and worry. They may react more strongly than others to stressful situations and struggle to control their aggression. In individuals with depression, aggression may be used as a way to alleviate pain and suffering. They may harm themselves and others to mask their pain and suffering. These individuals need professional support to manage their problems and find more constructive ways to cope with negative emotions. Finally,

in normal individuals, aggression may naturally occur in response to stressful situations. However, if aggression leads to physical or psychological harm, professional counseling and support are needed. Therefore, the treatment and management of aggression in individuals with anxiety, depression, and normal symptoms require different and specialized approaches. This involves a thorough understanding of the individual's history, analysis of symptoms, and identification of triggers and underlying causes of aggression to determine the best therapeutic strategies for each person.

On the other hand, the fear of loss was greater in the depression group compared to the anxiety and normal groups. Fear of loss is manifested by a strong desire to be aware of what others are doing. Some describe fear of loss as the fear of regret, leading to intense worry about missing out on social interactions, unique experiences, profitable investments, or other pleasurable events. Fear of loss can be defined from the perspective of self-determination theory as a state of poor self-monitoring, which forms due to a prolonged lack of real connection or difficulty in fulfilling psychological needs. In other words, fear of loss can be seen as intense anxiety about not being connected to social events, experiences, and interactions.

This finding can also be explained by stating that anxiety and depression crises can lead to fear of loss in individuals. Anxiety symptoms include constant worry, nervous tension, excessive anxiety, fear of unpleasant events occurring, and even physical symptoms. Depression symptoms include feelings of sadness and loss of appetite, self-disgust and dislike for others, decreased energy and interest in daily activities, and even suicidal thoughts. Treating fear of loss in individuals with anxiety and depression symptoms requires different approaches (Soheili-Pour et al., 2023). For individuals with anxiety symptoms, stress management techniques, breathing exercises, physical exercises, and even psychological counseling can be effective. For individuals with depression symptoms, psychological counseling, antidepressant medications, and even psychotherapy may be necessary. In any case, accurate diagnosis and timely treatment can help reduce fear of loss. Therefore, it is important for individuals with anxiety or depression symptoms to consult specialists and receive appropriate treatment.

5. Limitations & Suggestions

Despite the valuable results and new insights provided by this study, there are several limitations. The non-random sampling and use of convenience sampling methods may limit the generalizability of the results to a broader population. The use of self-report tools for data collection may involve response biases. Environmental factors, such as the setting for completing questionnaires and the daily stress and concerns of participants, may lead to changes in responses and affect the study results, compromising accuracy. Random sampling methods should be employed to enhance generalizability. Longitudinal methods for examining causal relationships between variables could be useful. To reduce response biases, combining self-report tools with more objective methods, such as semi-structured interviews or direct observation, is recommended. Intervention programs aimed at reducing fear of loss in individuals with anxiety and depression symptoms could significantly improve their quality of life. These programs could include psychological counseling and cognitive-behavioral therapies. Educating families and the community about the importance of recognizing and managing anxiety and depression symptoms could help reduce the psychological burden on these individuals. Workshops and educational courses for individuals with anxiety and depression symptoms and their families could be beneficial. Developing and using apps and online platforms to provide psychological counseling and support could increase access to psychological services and help reduce fear of loss. Further research into the relationship between fear of loss and other psychological factors could help identify new ways to reduce this fear.

Acknowledgments

We would like to express our appreciation and gratitude to all those who cooperated in carrying out this study.

Declaration of Interest

The authors of this article declared no conflict of interest.

Ethical Considerations

The study protocol adhered to the principles outlined in the Helsinki Declaration, which provides guidelines for ethical research involving human participants.

Transparency of Data

In accordance with the principles of transparency and open research, we declare that all data and materials used in this study are available upon request.

Funding

This research was carried out independently with personal funding and without the financial support of any governmental or private institution or organization.

Authors' Contributions

This article is derived from the first author's Master's thesis. All authors equally contributed in this article.

References

- Adavi, A., Fathi Marghmaleki, R., Madmoli, Y., & Madmoli, M. (2017). The Effectiveness of Group Cognitive-Behavioral Stress Management Interventions on Aggression in Women with Hypertension. <https://core.ac.uk/download/pdf/143844865.pdf>
- Aldahadha Al-Khawaldeh, B., & Mohammed Al-Khawalde, M. A. (2023). The Predictive Power of Ego Identity with Situational Regret Among the Divorced Women in Tafila Governorate. *Refereed Journal of Northern Europe Academy for Studies & Research (Denmark)*, 21. <https://ebsco-openurl.publicaciones.saludcastillayleon.es/results?sid=ebsco:ocu:record&bquery=DE+%22DIVORCED%20women%22>
- Bahmani Nia, M., & Sohrabi Shegefti, N. (2024). Comparison of Ego Strength, Defense Mechanisms, and Object Relations in Individuals with Depression and Healthy Individuals. *Clinical Psychology and Personality*, 22(1), 126-145. https://cpap.shahed.ac.ir/article_4316.html?lang=en
- Cheng, C. Y., & Wee, S. X. (2023). Kiasu (fear of losing out): An indigenous psychological construct in Singapore and its impact. *International Perspectives in Psychology: Research, Practice, Consultation*, 12(2), 65. <https://doi.org/10.1027/2157-3891/a000066>
- Cunningham, S., Hudson, C. C., & Harkness, K. (2021). Social media and depression symptoms: a meta-analysis. *Research on Child and Adolescent Psychopathology*, 49(2), 241-253. <https://doi.org/10.1007/s10802-020-00715-7>
- Daly, M., Sutin, A. R., & Robinson, E. (2021). Depression reported by US adults in 2017-2018 and March and April 2020. *Journal of affective disorders*, 278, 131-135. <https://doi.org/10.1016/j.jad.2020.09.065>
- Falla, D., Dueñas-Casado, C., & Ortega-Ruiz, R. (2023). Unjustified aggression in early childhood education: A systematic, narrative and conceptual review of the current scientific literature. *Aggression and Violent Behavior*. <https://doi.org/10.1016/j.avb.2023.101857>
- Hafer, A. (2020). *The Power of Sight: A Reexamination of Guercino and Poussin's Et in Arcadia Ego Paintings In Light of Early Modern Vision Theory* <https://etd.library.emory.edu/concern/etds/1c18dg89c>
- Harsanyi, S., Kupcova, I., Danisovic, L., & Klein, M. (2022). Selected biomarkers of depression: what are the effects of cytokines and inflammation? *International Journal of Molecular Sciences*, 24(1), 578. <https://doi.org/10.3390/ijms24010578>

- Hull, L., Levy, L., Lai, M. C., Petrides, K. V., Baron-Cohen, S., Allison, C., & Mandy, W. (2021). Is social camouflaging associated with anxiety and depression in autistic adults? *Molecular Autism*, 12, 1-13. <https://doi.org/10.1186/s13229-021-00421-1>
- Karami Boldaghi, R., & Daryazadeh, S. (2018). Predicting the elderly couple's decision-making styles based on adult attachment styles in Qeshm City [Research]. *Journal of Caspian Health and Aging*, 3(2), 14-23. <https://doi.org/10.22088/cjhaa.3.2.14>
- Kelly, J. R., Borre, Y., O'Brien, C., Patterson, E., El Aidy, S., Deane, J., & Hoban, A. E. (2016). Transferring the blues: depression-associated gut microbiota induces neurobehavioural changes in the rat. *Journal of psychiatric research*, 82, 109-118. <https://doi.org/10.1016/j.jpsychires.2016.07.019>
- Kinrade, C., Hart, W., & Lambert, J. T. (2022). Mental toughness, rather than self-esteem, moderates effects of grandiose narcissism on aggression under conditions of ego-threat. *Personality and individual differences*, 189, 111518. <https://doi.org/10.1016/j.paid.2022.111518>
- Kjærviik, S. L., & Bushman, B. J. (2021). The link between narcissism and aggression: A meta-analytic review. *Psychological bulletin*, 147(5), 477. <https://doi.org/10.1037/bul0000323>
- Madresi, S., & Shomali Askuei, A. (2023). Comparison of Ego Strength and Psychological Capital in Spouses of Recovered Addicts and Spouses of Non-Addicts. *Advances in Behavioral Sciences*, 8(56), 98-109. <https://ijndibs.com/article-1-810-fa.html>
- Malhotra, A., Kumari, S., & Koderma, J. A. (2023). Comparative Study of Ego Strength and Anxiety between Government and Private School Students of Ranchi Town. <https://www.scribd.com/document/660824437/A-Comparative-Study-of-Ego-Strength-and-Anxiety-between-Government-and-Private-School-Students-of-Ranchi-Town>
- Moncrieff, J., Cooper, R. E., Stockmann, T., Amendola, S., Hengartner, M. P., & Horowitz, M. A. (2023). The serotonin theory of depression: a systematic umbrella review of the evidence. *Molecular Psychiatry*, 28(8), 3243-3256. <https://doi.org/10.1038/s41380-022-01661-0>
- Nikosafat, Z., & Qarebaghi, F. (2020). The Mediating and Moderating Role of Ego Strength in the Relationship Between Object Relations and Identity States Among Students. *Clinical Psychology and Personality (Daneshvar Raftar)*, 18(2), 109-118. <https://www.sid.ir/paper/1044139/en>
- Radomsky, A. S. (2022). The fear of losing control. *Journal of Behavior Therapy and Experimental Psychiatry*, 77, 101768. <https://doi.org/10.1016/j.jbtep.2022.101768>
- Ramazani, K., & Hadizadeh Kafash, R. (2016). Effectiveness of Hypnotherapy on Aggression in Students.
- Rapee, R. M., Creswell, C., Kendall, P. C., Pine, D. S., & Waters, A. M. (2023). Anxiety disorders in children and adolescents: A summary and overview of the literature. *Behaviour Research and Therapy*. <https://doi.org/10.1016/j.brat.2023.104376>
- Renaud-Charest, O., Lui, L. M., Eskander, S., Ceban, F., Ho, R., Di Vincenzo, J. D., & McIntyre, R. S. (2021). Onset and frequency of depression in post-COVID-19 syndrome: A systematic review. *Journal of psychiatric research*, 144, 129-137. <https://doi.org/10.1016/j.jpsychires.2021.09.054>
- Sadafi, E., Monirpour, N., & Mirzahoseini, H. (2020). Explaining Structural Relationships of the Relationship between Objective Relationship Quality and Marital Instability Based on the Mediating Role of Ego Power. *medical journal of mashhad university of medical sciences*, 63(Special Psychology). https://mjms.mums.ac.ir/article_20771.html
- Soheili-Pour, A., Rashtbari, A., Ahmadi, D., & Saed, O. (2023). Changes in Symptoms of Generalized Anxiety, Depression, Obsessive-Compulsive Disorder, Social Anxiety, and Generalized Anxiety Among Students During the COVID-19 Pandemic. *Cognitive Psychology and Psychiatry*, 10(1), 45-58. <https://doi.org/10.32598/shenakht.10.1.45>
- Tam, K. P., Chan, H. W., & Clayton, S. (2023). Climate change anxiety in China, India, Japan, and the United States. *Journal of Environmental Psychology*, 87, 101991. <https://doi.org/10.1016/j.jenvp.2023.101991>
- Tao, Y., Tang, Q., Zou, X., Wang, S., Ma, Z., Zhang, L., & Liu, X. (2023). Effects of attention to negative information on the bidirectional relationship between fear of missing out (FoMO), depression and smartphone addiction among secondary school students: Evidence from a two-wave moderation network analysis. *Computers in human Behavior*, 148, 107920. <https://doi.org/10.1016/j.chb.2023.107920>
- Toprak, M., Boldrini, C., Passarella, A., & Conti, M. (2022). Harnessing the power of ego network layers for link prediction in online social networks. *IEEE Transactions on Computational Social Systems*, 10(1), 48-60. <https://doi.org/10.1109/TCSS.2022.3155946>
- Wei, D., Osakada, T., Guo, Z., Yamaguchi, T., Varshneya, A., Yan, R., & Lin, D. (2023). A hypothalamic pathway that suppresses aggression toward superior opponents. *Nature Neuroscience*, 26(5), 774-787. <https://doi.org/10.1038/s41593-023-01297-5>
- Wei, L., Yang, Y., Wu, J., Long, C., & Li, B. (2022). Trust management for Internet of Things: A comprehensive study. *Ieee Internet of Things Journal*, 9(10), 7664-7679. <https://doi.org/10.1109/JIOT.2021.3139989>
- Wisse, B., Rus, D., Keller, A. C., & Sleebos, E. (2019). "Fear of losing power corrupts those who wield it": the combined effects of leader fear of losing power and competitive climate on leader self-serving behavior. *European Journal of work and organizational psychology*, 28(6), 742-755. <https://doi.org/10.1080/1359432X.2019.1635584>
- Zanjanchi Niko, S., & Farahani, A. (2024). Comparison of Ego Strength and Defense Mechanisms in Patients with Borderline Personality Disorder and Healthy Individuals. *Khomeini Shahr*.